



## MULTICOMPONENT SYNTHESIS AND EVALUATION OF ANTIBACTERIAL ACTIVITY OF BENZOTHIAZOLO PYRIMIDO-PYRIMIDO BENZOTHIAZOLE AND ITS DERIVATIVES

Anil Chidrawar\*

\*Research Centre & P.G. Department of Chemistry, Degloor College, Degloor, Pin- 431717. S.R.T.M.U. Nanded, Maharashtra (India)

\*Corresponding Author Email: [anilchidrawar74@gmail.com](mailto:anilchidrawar74@gmail.com)

### ABSTRACT

One-pot multicomponent reactions constitute an especially attractive recent synthetic strategy since they provide easy and rapid access to a large number of organic compounds with diverse substitution pattern with short time. To synthesize fused condensed pyrimido benzothiazole possessing more than three rings which exhibit a wide spectrum of biological and pharmacological activities. We report the synthesis of 14, 15-diimino-10-nitro-benzothiazolo [2,3-b] pyrimido [5,6-e] pyrimido [2,1-b] [1,3] benzothiazole with condensation of 3-cyano-4-imino-2-methylthio-8-nitro-4H-pyrimido [2,1-b] [1,3] benzothiazole and 2-amino-6-substituted benzothiazole in the presence of DMF and catalytic amount of  $K_2CO_3$  by refluxing 5-6 hours by one pot multicomponent synthesis. In conclusion a facile one pot synthesis has been developed for the title compounds using readily available starting materials. All the newly synthesized compounds were screened for antibacterial activity. These compounds were found to possess a broad-spectrum activity.

### KEY WORDS

2-amino-6-substituted benzothiazole, DMF, Potassium carbonate, Pyrimido benzothiazole

### INTRODUCTION

A survey of literature made it evident that, very little work has been carried out on the synthesis of fused pyrimido benzothiazole possessing three to four rings which exhibit a wide spectrum of biological and pharmacological activities like anti-allergic, antiparkinsonium[1], herbicidal[2], antiviral, phosphodiesterase inhibition, anti-parasitic activity [3], anti-inflammatory [4] and antitumor activity.

One pot synthesis is a green approach towards the synthesis of various heterocyclic compounds and for a researcher there is lot of scope to change the reaction condition, to change the catalyst, to change the solvent or to modify the catalyst. Pyrimidine, iminopyrimidine, pyrazole and fused benzothiazole heterocycles are reported to be effective pharmacophores [5-11]. Nair


Mohan D. *et.al.* synthesized pyrimido [2,1-b] benzothiazole and its derivatives by refluxing diethyl ethoxy methyle nemalonate with respective 2-amino benzothiazole found to be antiviral activity.

A comprehensive review on the methods for the synthesis of iminopyrimidines has been published in the form of book, "The Pyrimidines" by Brown D.J. *et al.* [12-13]. Imino compounds are known to possess some sedative and hypnotic actions. Denny W.A. *et al.* [14] reported anticancer agents from fused pyrido-imidazo, -pyrazolo, -pyrazino and pyrrolo heterocycles. Jimonet Patrick and his research group [15] reported synthesis of pharmacologically active 6-(trifluoromethoxy)- 3-substituted-2-imino benzothiazolines. Erlenmeyer and Von Meyenburg [16] reported moderate sedative activity of 5, 5-dialkyl-2-imino-4-thiazolidones which is

in marked contrast to the lack of activity of the iminobarbituric acids [17-20]. In view of these reported biological activities of this system, synthesis of such condensed system has attracted much attention in recent years. In this note, we report one pot multicomponent synthesis of 14, 15-diimino-10-nitro-benzothiazolo [2,3-*b*] pyrimido [5,6-*e*] pyrimido [2,1-*b*] [1,3] benzothiazole and its derivatives.

## MATERIAL AND METHODS

**Experimental:** All melting points were determined in open capillary tube and were uncorrected. IR spectra were recorded with potassium bromide pellets technique,  $^1\text{H}$  NMR spectra were recorded on AVANCE 300 MHz Spectrometer in DMSO using TMS as internal standard. Mass spectra were recorded on a FT VG-7070 H Mass Spectrometer using EI technique at 70 eV. All the reactions were monitored by Thin layer chromatography.



### General procedure

#### Multicomponent Synthesis of 3-substituted derivatives of 14,15-diimino-10-nitro-benzothiazolo [2,3-*b*] pyrimido [5,6-*e*] pyrimido [2,1-*b*] [1,3] benzothiazole (C1-C5).

It is prepared by condensation of 3-cyano-4-imino-2-methylthio-8-nitro-4H-pyrimido [2,1-*b*] [1,3] benzothiazole (A) independently with 2-amino benzothiazole (B1), 2-amino-6-methyl benzothiazole (B2), 2-amino-6-methoxy benzothiazole (B3), 2-amino-6-chloro benzothiazole (B4), and 2-amino-6-nitro benzothiazole (B5) in the presence of DMF and catalytic amount of K<sub>2</sub>CO<sub>3</sub> reflux for 5-6 hours to get 14,15-diimino-10-nitro-benzothiazolo[2,3-*b*] pyrimido [5,6-*e*] pyrimido [2,1-*b*] [1,3] benzothiazole and its 3-substituted derivatives (C1-C5).

#### Multicomponent Synthesis of 14,15-diimino-10-nitro-benzothiazolo [2,3-*b*] pyrimido [5,6-*e*] pyrimido [2,1-*b*] [1,3] benzothiazole (C1).

Condensation of 3-cyano-4-imino-2-methylthio-8-nitro-4H-pyrimido[2,1-*b*] [1,3] benzothiazole (A) reflux with 2-amino benzothiazole (B1) in the presence of DMF and

catalytic amount of K<sub>2</sub>CO<sub>3</sub> for 5-6 hours to yields 14,15-diimino-10-nitro-benzothiazolo[2,3-*b*] pyrimido[5,6-*e*] pyrimido [2,1-*b*] [1,3] benzothiazole.

Yield : 59 %, IR:(KBr/cm<sup>-1</sup>) : 3281 & 3322 (=NH), 3111 (Ar-H), 1615 (C=N), 1517 & 1350 (NO<sub>2</sub>),  $^1\text{H}$ -NMR: (DMSO):  $\delta$  4.01 (s 1H =NH),  $\delta$  4.11 (s 1H =NH),  $\delta$  7.10 (d 4H Ar-H),  $\delta$  7.11 (d 3H Ar-H), EI-MS: (m/z:RA%): 420 (M+1), Elemental analysis : C<sub>18</sub>H<sub>9</sub>N<sub>7</sub>O<sub>2</sub>S<sub>2</sub>, Calculated: (%) C 51.54, H 2.16, N 23.28, O 7.63, S 15.29 Found (%) : C 51.51, H 2.11, N 23.20, O 7.60, S 15.24.

#### Multicomponent Synthesis of 3-methyl-14,15-diimino-10-nitro-benzothiazolo[2,3-*b*] pyrimido [5,6-*e*] pyrimido [2,1-*b*] [1,3] benzothiazole (C2).

Condensation of 3-cyano-4-imino-2-methylthio-8-nitro-4H-pyrimido[2,1-*b*] [1,3] benzothiazole (A) and 2-amino-6-methyl benzothiazole (B2) in the presence of DMF and small amount of potassium carbonate for 5 hours to give 3-methyl-14,15-diimino-10-nitro-benzothiazolo[2,3-*b*] pyrimido[5,6-*e*] pyrimido[2,1-*b*] [1, 3] benzothiazole.

Yield : 55 %, IR:(KBr/cm<sup>-1</sup>) : 3310 & 3325 (=NH), 3110 (Ar-H), 1616 (C=N), 1522 & 1340 (NO<sub>2</sub>), EI-MS: (m/z:RA%): 434 (M+1), Elemental analysis : C<sub>19</sub>H<sub>11</sub>N<sub>7</sub>O<sub>2</sub>S<sub>2</sub>, Calculated: (%) C 52.65, H 2.56, N 22.62, O 7.38, S 14.79 Found (%) : C 52.61, H 2.50, N 22.60, O 7.31, S 14.75.

**Multicomponent Synthesis of 3-methoxy-14, 15-diimino-10-nitro-benzothiazolo [2, 3-b] pyrimido [5, 6-e] pyrimido [2, 1-b] [1, 3] benzothiazole (C3).**

Condensation of 3-cyano-4-imino-2-methylthio-8-nitro-4H-pyrimido[2,1-b] [1,3] benzothiazole (A) and 2-amino-6-methoxy benzothiazole (B3) in the presence of Dimethyl formamide and potassium carbonate for 6 hours to get 3-methoxy-14,15-diimino-10-nitro-benzothiazolo[2,3-b] pyrimido[5,6-e] pyrimido[2,1-b] [1,3] benzothiazole.

Yield : 57 %, IR:(KBr/cm<sup>-1</sup>) : 3312 & 3313 (=NH), 3100 (Ar-H), 1620 (C=N), 1515 & 1332 (NO<sub>2</sub>), 1160 (C-O), EI-MS: (m/z:RA%): 450 (M+1), Elemental analysis : C<sub>19</sub>H<sub>11</sub>N<sub>7</sub>O<sub>3</sub>S<sub>2</sub>, Calculated: (%) C 50.77, H 2.47, N 21.81, O 10.68, S 14.27 Found (%) : C 50.70, H 2.44, N 21.76, O 10.62, S 14.25

**Multicomponent Synthesis of 3-chloro-14,15-diimino-10-nitro-benzothiazolo[2,3-b] pyrimido [5,6-e] pyrimido [2,1-b] [1,3] benzothiazole (C4).**

Condensation of 3-cyano-4-imino-2-methylthio-8-nitro-4H-pyrimido[2,1-b] [1,3] benzothiazole (A) and 2-

amino-6-chloro benzothiazole (B4) in the presence of Dimethyl formamide and potassium carbonate for 5 hours to produce 3-chloro-14,15-diimino-10-nitro-benzothiazolo [2, 3-b] pyrimido [5, 6-e] pyrimido[2,1-b] [1,3] benzothiazole.

Yield : 70 %, IR:(KBr/cm<sup>-1</sup>) : 3312 & 3280 (=NH), 3111 (Ar-H), 1627 (C=N), 1519 & 1335 (NO<sub>2</sub>), EI-MS: (m/z:RA%): 454 (M+1), Elemental analysis: C<sub>18</sub>H<sub>8</sub>ClN<sub>7</sub>O<sub>2</sub>S<sub>2</sub>, Calculated: (%) C 47.63, H 1.78, Cl 7.81, N 21.60, O 07.05, S 14.13 Found (%) : C 47.61, H 1.74, Cl 7.75, N 21.55, O 07.00, S 14.10

**Multicomponent Synthesis of 3-nitro-14, 15-diimino-10-nitro-benzothiazolo [2, 3-b] pyrimido [5,6-e] pyrimido [2,1-b] [1,3] benzothiazole (C5).**

Condensation of 3-cyano-4-imino-2-methylthio-8-nitro-4H-pyrimido[2,1-b] [1,3] benzothiazole (A) and 2-amino-6-nitro benzothiazole (B5) in presence of DMF and K<sub>2</sub>CO<sub>3</sub> for 6 hours to produce 3-nitro-14,15-diimino-10-nitro-benzothiazolo [2,3-b] pyrimido[5,6-e] pyrimido [2,1-b] [1,3] benzothiazole.

Yield : 64 %, IR:(KBr/cm<sup>-1</sup>) : 3300 & 3265 (=NH), 3121 (Ar-H), 1622 (C=N), 1511 & 1355 (NO<sub>2</sub>), EI-MS: (m/z:RA%): 465 (M+1), Elemental analysis : C<sub>18</sub>H<sub>8</sub>N<sub>8</sub>O<sub>4</sub>S<sub>2</sub>, Calculated: (%) C 46.55, H 1.74, N 24.13, O 13.78, S 13.81 Found (%) : C 46.51, H 1.70, N 24.10, O 13.74, S 13.74

**Table-1: Antibacterial activity of 3-substituted derivatives of 14, 15-diimino-10-nitro-benzothiazolo [2,3-b] pyrimido [5,6-e] pyrimido [2,1-b] [1,3] benzothiazole (C1-C5).**

Sample code	*Zone of inhibition (diameter in mm)			
	<i>B. subtilis</i>	<i>B. Megatenium</i>	<i>E. coli</i>	<i>P. aureginosa</i>
	100µ/ml	100µ/ml	100µ/ml	100µ/ml
C1	21	26	07	10
C2	27	23	10	09
C3	13	15	20	22
C4	10	09	25	19
C5	23	20	10	11
Streptomycin	31	35	28	27
DMSO	-	-	-	-

\*Each value is an average of three independent determinations ± Standard deviation.

**Note:** '-' denotes no activity, 7-12 mm poor activity, 13-17 mm moderate activity, 18-27 mm and above good activity.

**RESULT AND DISCUSSION**

One pot reaction constitutes an especially attractive recent synthetic strategy since they provide easy and rapid access to a large number of organic compounds with diverse substitution pattern. In the present work, we report one-pot synthesis of a novel fused

heterocyclic compound, 14,15-diimino-10-nitro-benzothiazolo [2,3-b] pyrimido [5,6-e] pyrimido [2,1-b] [1,3] benzothiazole and its 3-substituted derivatives (C1-C5). All newly synthesized derivatives (C1-C5) were evaluated in-vitro for antibacterial activity against gram positive and gram-negative bacterial strain such as

*Bacillus subtilis*, *Bacillus Megatenium*, *Escherichia coli* and *Pseudomonas aureginosa* at concentration 100 $\mu$ /ml by disc diffusion method<sup>24</sup> by using DMSO as solvent control and nutrient agar was employed as culture media. After 24h of incubation at 37<sup>o</sup>C, the zone of inhibition were measured in mm. The activity was compared with known antibiotic Streptomycin and the data was represented in Table-1.

### CONCLUSION

In conclusion a facile one pot synthesis has been developed for the title compounds using readily available starting materials. All the newly synthesized compounds were screened for antibacterial activity. These compounds were found to possess a broad-spectrum activity. However, the activities of the tested compounds are much less than those of standard antibacterial agents used. All newly synthesized compounds were screened for antibacterial activity studies at a concentration of 100 $\mu$ /ml using DMSO as a control and Streptomycin used as a standard against gram positive and gram-negative bacteria. The data in Table 1 indicates that compound C1, C2 and C5 was found to possess a broad-spectrum activity against gram-positive bacteria and compound C3 and C4 were found to possess a broad-spectrum activity against gram-negative bacteria. However, the activities of the tested compounds are less than those of standard antibacterial agents used.

### ACKNOWLEDGEMENTS

The authors are thankful to the Director, Indian Institute of Chemical Technology, Hyderabad for providing spectra.

### CONFLICTS OF INTERESTS

Author has none to declare.

### REFERENCES

- [1] Wade JJ, Toso CB, Matson CJ, Stelzer VL. Synthesis and antiallergic activity of some acidic derivatives of 4H-pyrimido [2,1-b] benzazol-4-ones. *J Med Chem*, 26(4): 608-11, (1983)
- [2] Vishnuji Ram. Synthesis of Pyrimidines and Fused Pyrimidines as leishmanicides and herbicides. *J Prakt Chem*, 331(6): 893-905, (1989)
- [3] Alaimo RJ. The synthesis of some 4H-pyrimido[2,1-b] benzothiazol-4-ones. *J Het Chem*, 10(5): 769-72 (1973)
- [4] Glennon RA, Gaines JJ, Rogers ME. Benz-fused mesoionic xanthine analogs as inhibitors of cyclic-AMP phosphodiesterase. *J Med Chem*, 24(6): 766-9, (1981)
- [5] Bartovic Alexander, Dusas Ilavsky, Ondrej Simo, Zalibera, Belicova Lubomir, Anna, Milan Seman. Synthesis of Nitro-Substituted 4-Oxo-4H-pyrimido[2,1-b] benzothiazole-3-carboxylic Acids and their Spectral Characteristics. *Collection Czech Chem Commun*, 60(4): 583-93, (1995)
- [6] Dunwell DW and D Evans. The reactions of 2-aminothiazoles and 2-aminobenzothiazoles with propiolic acid and its esters. *J Chem Soc*, 0: 2094-2097, (1971)
- [7] Hataba AA, Fikry RM, Moustafa HY. Synthesis of New Benzothiazole And Condensed Benzothiazole Derivatives. *J Ind Chem Soc*, 74: 818-819, (1997)
- [8] Reimlinger H, Peiren MA, Merenyi R. *Chem Ber Synthesen mit heterocyclischen Aminen*, 108(12): 3894-97, (1975)
- [9] Sakamoto Masanori, Miyazawa Kyoko, Tomimatsu Yoshid. Acids & vitamin B<sub>1</sub> have remarkable pharmaceutical importance by the reaction of dimethyl allene-1,3-dicarboxylate. *Chem Pharm Bull*, 25(12): 3360-3365, (1977)
- [10] Kutyzev Alexander, Kappe Thomas. Benzimidazole condensed ring systems. *J Het Chem*, 36(11): 237-240, (1999)
- [11] Fatemeh Chadegani, Fatemeh Darviche, Saeed Balalaie. A New and Efficient Method for the Synthesis of Pyrimido[2,1-b] benzothiazole derivatives. *International Journal of Organic Chemistry*, 2(1): 31-37, (2012)
- [12] Deshmukh Vinayak K, Yadav Mukesh R, Chaudhari Sanjay. Microwave Assisted Synthesis and Anticancer Activity of Substituted Pyrimido [2,1-b] [1,3] benzothiazole derivatives. *Int J Pharm Sci Rev Res*, 22(1): 41-47, (2013)
- [13] "The Pyrimidines" Brown DJ, Evans RF, Cowden WB and Feun MD. John Wiley and Sons, New York, 1993.
- [14] Denny WA, Gamage SA, Spicer JA, Rewcastle GW, Milton J, Sohail S, Dangerfield W, Mistry P, Vicker N, Charlton PA, Structure-activity relationships for pyrido, imidazo, pyrazolo, pyrazino and pyrrolophenazine carboxamides as topoisomerase targeted anticancer agents. *J Med Chem*, 45(3): 740-743, (2002)
- [15] Erlenmeyer and Von Meyenburg. Synthesis of pharmacologically active 6-(trifluoromethoxy)- 3-substituted-2-imino benzothiazolines. *Chim Acta*, 20: 1388, (1937)
- [16] Rana Arpana, Siddiqui N, Khan SA. Benzothiazoles: A new profile of biological activity. *Ind J Pharm Sci*, 69(1): 10-17, (2007)
- [17] Kini S, Swain SP and Gandhi AM. Synthesis and evaluation of novel benzothiazole derivatives against human cervical cancer cell lines. *J Pharm Sci*, 69(1): 46-50, (2007)
- [18] TD Bradshaw, DF Shi, RJ Schultz, KD Paull, L Kelland, A Wilson, HH Fiebig, S Wrigley and MFG Stevens. Influence of 2-(4-aminophenyl) benzothiazoles on growth of human



ovarian carcinoma cells in vitro and in vivo. *Br J Cancer*, 78: 421-429, (1998)

[19] I Hutchinson, M Chua, HI Browne, V Trapani, TD Bradshaw, AD Westwell and MFG Stevens. Synthesis and in vitro biological properties of fluorinated 2-(4-

aminophenyl) benzothiazoles. *J Med Chem*, 44(9): 1446-55, (2001)

[20] I Hutchinson, SA Jennings, BR Vishnuvajjala, AD Westwell and MFG Stevens. Synthesis and pharmaceutical properties of antitumor 2-(4-aminophenyl) benzothiazole amino acid prodrugs. *J Med Chem*, 45(3), 744-47 (2002)

Received:05.05.18, Accepted: 08.06.18, Published:01.07.2018

**\*Corresponding Author:**

**Anil Chidrawar\***

Email: [anilchidrawar74@gmail.com](mailto:anilchidrawar74@gmail.com)